

# Digestive Dilemmas

An overview of GI issues in horses



ABIGAIL BOATWRIGHT

LUCILE VIGOUROUX

**H**orses are known for having delicate gastrointestinal systems. Equine digestion is full of twists and turns, and a lot can go wrong during the process. Colic, diarrhea, ulcers, and other ailments are common in our domestic horse population. Being informed and prepared will help you react appropriately should your horse ever greet you with a bellyache.

## **Colic: The Main Culprit**

Colic is the No. 1 killer of adult horses, accounting for 30% of U.S. horse deaths, according to the USDA's 2015 National Animal Health Monitoring System study. While the clinical signs of colic—an umbrella term referring to abdominal discomfort—might be similar in many cases, the underlying causes and conditions can be vastly different and have a major influence on treatment and prognosis.

Michael Fugaro, VMD, Dipl. ACVS, owner of

Mountain Pointe Equine Veterinary Services and a surgeon at B.W. Furlong and Associates, both in New Jersey, says the two most common types of colic he encounters on the farm are gas/spasmodic (gut spasms) and idiopathic (unknown cause), with impaction next on the list.

“Thankfully, most of these cases respond to a single treatment of medical therapy and resolve without further complication,” he says, for example, administering a non-steroidal anti-inflammatory

drug (NSAID) such as flunixin meglumine (Banamine) under your vet's guidance.

So what causes colic? "Horses are creatures of habit," Fugaro says, "and when changes in feed, grass, water intake, medication, exercise regime, stabling, environment, etc. occur, the intestinal tract appears to be the most susceptible to that impact."

Types of colic that end up requiring surgery vary depending on the region and horse demographic. This is because different horses in different parts of the country are exposed to different risk factors. "The signalment (age, sex, breed), occupation, and other variables of the horse appear to influence the type of surgical colic experienced," he says. "For instance, older horses and Arabians are highly represented with strangulating obstructions secondary to lipomas (benign fatty tumors); geldings that crib-bite and/or windsuck have a higher incidence of entrapment of the small intestine in the epiploic foramen (a narrow opening

connecting the two sacs of the abdominal cavity); and broodmares show a higher prevalence of large colon (the stretch of intestine located between the cecum and the transverse colon) displacement and/or volvulus (twisting) after foaling."

Nimet Browne, DVM, MPH, Dipl. ACVIM, an internal medicine specialist at Hagyard Equine Medical Institute, in Lexington, Kentucky, with an interest in gastrointestinal disease, echoes these observations. Located in the heart of Kentucky horse country, Browne treats many Thoroughbred broodmares and foals. Common causes of medical colic in her patient population include large colon displacements, spasmodic colic, enteritis (inflammation of the small intestine), and impaction, including from parasites.

The colic type a horse experiences also depends on his age and lifestyle. "The majority of the colic cases referred to us are medical colics (those that don't need surgery because they can be managed with intravenous fluids, pain-relieving

medications, and supportive care)," Browne says. "A colic case can require surgery if pain cannot be controlled, if there is indication of gastrointestinal compromise, or if all other treatment options have failed."

Researchers have confirmed that the faster a horse that isn't responding to medical therapy gets referred to surgery, the greater his chance of a successful outcome. "Surgical causes of colic that we frequently see include large colon volvulus, small intestinal volvulus, mesenteric rents (tears in the membrane that anchors the intestines to the abdominal wall), intussusceptions (where the intestine telescopes onto itself), and colonic displacements," Browne says.

While it can be difficult for veterinarians to determine the cause of a colic, she sees certain well-known predisposing factors in her practice population, she adds. These include the immediate peripartum (post-birthing) period, advanced age, stress from yearling sales prepping,



# GutWerks®

**Probiotics, Prebiotics, Active Yeast  
plus Yeast Culture**

GutWerks offers a super-concentrated boost of our popular probiotic/prebiotic package along with premium quality yeast culture and live cell yeast! Our probiotics, prebiotics and live yeast cultures have been selected for performance, viability and stability when mixed in feed—they'll be there when you need them.

Each two (2) ounce serving of GutWerks will provide over 20 billion CFU of live, naturally-occurring, beneficial microorganisms and 50 billion live *Saccharomyces cerevisiae* yeast cells. GutWerks also includes our prebiotic package (mannanoligosaccharides and fructooligosaccharides), all on a super-palatable Diamond V yeast culture base.

## HorseTech®

Awesome Supplements...  
Amazing Service!

Always Free  
Shipping!

QUALITY SEAL



NATIONAL ANIMAL SUPPLEMENT COUNCIL

Order Online at: [www.horsetech.com](http://www.horsetech.com)

Order by Phone at: (800) 831-3309    Email: [service@horsetech.com](mailto:service@horsetech.com)    Free Shipping in Contiguous U.S.





horse shows, stall rest, or, as mentioned, intestinal parasite load.

**Diarrhea: How Serious Can it Be?**

In short, very. Although occasional loose manure can simply reflect recent feed changes or a brief stressful episode, severe diarrhea (due to toxins, parasites, infectious agents such as *Salmonella*, and more) can signify something is seriously wrong.

“Diarrhea cases are not to be taken lightly and can become fatal in severe cases,” says Fugaro. “If a horse has true diarrhea (waterlike manure that immediately sinks below the bedding), it is considered a medical emergency.”

Here’s why: Diarrhea can cause severe dehydration and protein loss, and advanced cases can even lead to the debilitating hoof condition laminitis and bodywide infection in the blood known as sepsis. In addition, acute diarrhea is very difficult to treat.

“Many horses will require around-the-clock aggressive medical and supportive therapies that can only be provided in a hospital setting,” says Fugaro.

So, how can you gauge the severity of a horse’s diarrhea? “It is of utmost importance to consider the systemic health of the horse in question,” Browne says. “Mild diarrhea caused by feed changes is typically transient (one to three days at the most) and not generally accompanied



COURTESY AMY MCLEAN

A colic case can require hospitalization and surgery if pain cannot be controlled, if there is indication of gastrointestinal compromise, or if all other treatment options have failed.

by changes in attitude, demeanor, appetite, or body temperature.”

If any of the latter signs do appear, she urges owners to call their veterinarians and ensure their horses have plenty of drinking water available, temporarily remove concentrates from the diet, and potentially administer a veterinarian-

prescribed adsorbent to help neutralize the diarrhea-causing toxins. She also recommends asking your veterinarian whether you should give your horses probiotics (beneficial microorganisms) and prebiotics (micronutrients the good gut microbiota require to survive). Further care, such as antimicrobials and even hospitalization, might be necessary.

**IS YOUR HORSE'S HEALTH LEAKING AWAY?** KEMIN EQUINE

Leaks in your horse's intestinal barrier — known as **Leaky Gut Syndrome** — cause intestinal inflammation and can overflow into various health and welfare problems. Good nutrition and attention to gut health can alleviate the negative impacts of stress and help promote:

- Immunity
- Performance
- Healthy weight
- Nutrient absorption
- Positive attitude

kemin.com/leakygut GUT HEALTH SOLUTION

Ensure your feed includes **CLOSTAT®** and **ButiPEARL® Z EQ** today.

© Kemin Industries, Inc. and its group of companies 2020. All rights reserved. \*\*The trademarks of Kemin Industries, Inc. U.S.A.

**Ulcers: A Pain in the Gut**

Equine gastric ulcer syndrome (EGUS), the most common gastrointestinal disease in horses, refers to ulcerative lesions of the stomach. Equine colonic ulcer syndrome (ECUS) is a similar condition that affects the horse’s hindgut—the digestive tract beyond the small intestine. Ulcers cause discomfort, pain, and often a decrease in body condition. Other reported effects include poor appetite, attitude changes, lack of energy, a dull hair coat, and chronic diarrhea.

While they might not be immediately evident, ulcers are prevalent among pleasure, performance, and especially racehorse populations and linked to many factors. Fasting, lack of forage, high amounts of concentrate feed, stress, stall confinement, transport, and NSAID administration are among the most

common. However, there are many steps you can take to help protect your horse's stomach, starting by providing as much turnout and forage access as possible. If your horse needs NSAIDs to manage pain associated with arthritis or other chronic conditions, for instance, ask your veterinarian about using firocoxib (Equioxx), a COX-2 selective NSAID that's gentler on the stomach than nonselective forms like Bute or Banamine.

Omeprazole, an oral antiacid sold under the names GastroGard and UlcerGard, is highly effective in treating and preventing gastric ulcers. Colonic ulcers are more difficult to diagnose and eradicate but have anecdotally responded to gastroprotectants such as misoprostol.

Consult your veterinarian if you suspect your horse has ulcers. He or she might perform gastroscopy (passing an endoscope through the nostrils and into the stomach to look at its surface) to make a diagnosis and can prescribe the appropriate treatment.

### Gut Microbiome Disturbances

The horse's intestines are populated with beneficial bacteria, protozoa, fungi, and archaea (single-celled organisms) responsible for digesting and absorbing nutrients. Together, they make up the microbiome, which digests fiber into volatile fatty acids the body can use for

energy. Any alteration of the flora within the digestive tract, known as dysbiosis, can impair the gut's normal functioning, potentially causing big problems.

"The microbiome is highly influenced by diet, geographic location, and host genetics, as well as systemic medications and/or supplements," Browne says.

“General principles of restoring a healthy microbiome include removing inciting causes, administering probiotics and prebiotics, and providing high-quality fiber.”

DR. NIMET BROWNE

"When the microbiome is altered, minor to life-threatening complications can arise, including colic, colitis (inflammation of the large colon), laminitis, gastric ulcers, as well as weight loss, obesity, diarrhea, and systemic inflammation."

When treating these conditions, she stresses the importance of identifying and addressing underlying problems, which could include inflammatory bowel

conditions and/or infectious causes of diarrhea.

"Even once these issues are addressed, maintaining and cultivating normal flora can be difficult," Browne says. "General principles of restoring a healthy microbiome include removing inciting causes, administering probiotics and prebiotics, and providing high-quality fiber."

### Malabsorption of Nutrients

Malabsorption occurs when the intestinal mucosa fails to absorb and transport nutrients such as carbohydrates, protein, fat, vitamins, or minerals into the bloodstream correctly. "There is ongoing research about malabsorption, as it is believed to be a contributing factor in many of the clinical symptoms we observe in horses," Fugaro says.

Browne adds that malabsorption usually occurs when inflammatory or neoplastic (abnormal tissue growth) cells infiltrate the intestinal mucosa. Inflammation of the small intestine usually leads to chronic weight and protein loss, while that of the large intestine (colitis) more commonly causes diarrhea. Other clinical signs include lethargy, colic, and dependent edema (swelling affected by gravity, generally on the abdomen and/or legs).

"Chronic inflammatory bowel disease, which is commonly diagnosed in humans but more rarely in horses, is often a culprit," Browne says. "So are neoplasia, small strongyles (intestinal parasites), diffuse fibrosis (scarring), extensive small intestinal resection during colic surgery, and also infectious conditions such as (equine proliferative enteropathy)."

Again, treatment depends on the cause of the malabsorption, which is best determined by your veterinarian.

### Take-Home Message

Horses have sensitive gastrointestinal tracts that are susceptible to a variety of ailments, but owners can take several simple steps to optimize digestive health. Provide constant access to forage and fresh water; make feed changes gradually, turn your horse out as much as possible, and reduce sources of stress in his environment. And, as always, communicate early signs of trouble to your veterinarian. With gastrointestinal issues, rapid intervention is crucial for a successful outcome. 🐾

Available through your Veterinarian